

Investigation of Super Lehmer-3 mean Labeling in Theta Related Graphs

D. Praveen Prabhu^{1,2,*}, O. V. Shanmuga Sundaram¹

¹Department of PG Mathematics, Sree Saraswathi Thiagaraja College, Thippampatti, Pollachi, Coimbatore - 642 107, Tamil Nadu, India

²Department of Mathematics, PPG Institute of Technology, Coimbatore - 641 035, Tamil Nadu, India

Abstract In graph theory, labeling of vertices and edges according to specific rules offers insight into the structural properties of graphs and their potential applications. This study focuses on a particular type of graph labeling known as Super Lehmer-3 Mean Labeling. A graph is said to admit such a labeling if there exists an injective vertex labeling such that the corresponding edge labels, derived from the Lehmer-3 mean of their incident vertex labels, are also distinct and satisfy certain arithmetic conditions. Graphs that admit such a labeling are termed Super Lehmer-3 Mean Graphs. In this work, we investigate the existence of super Lehmer-3 mean labeling in various families of theta-related graphs. Specifically, we analyze standard theta graphs and their structural variations obtained by subdivisions and attachments of pendant vertices. The results confirm that these modified forms of theta graphs also admit super Lehmer-3 mean labeling. Additionally, we extend our exploration to harmonic mean labeling in selected theta graphs, highlighting the conditions under which this alternative labeling scheme is valid. The findings presented in this paper contribute to the growing body of knowledge in graph labeling theory, offering new characterizations and constructions of labeled graph families. These results may have implications for theoretical studies and applications where such labeling structures are relevant.

Keywords Graph, Lehmer-3 Mean Labeling, Super Lehmer-3 Mean labeling, Harmonic Mean Labeling, Corona Product of G , Subdivision of G , Theta Graph

DOI: 10.19139/soic-2310-5070-2633

1. Introduction

“Graph Theory” is a very important branch of arithmetic, Euler (1707-1782) is understood because the father of Graph Theory likewise as Topology. Graph Theory came into existence throughout the primary half the 18th century. Graph Theory failed to begin to become associate organized branch of mathematics till the last half of the 19th century and, there wasn't even a book on the topic till the primary half the 20th century. Graph Theory is a branch of mathematics concerned with networks of points connected by lines. The subject of graph theory had its beginning in recreational with problems but it has grown into a significant area of mathematical research, application in chemistry, operations research, social science, application in biology and computer science.

Labeled graphs provide mathematical models for a broad range of applications. The qualitative labeling of graph elements have been used in diverse fields such as conflict resolutions in social psychology, energy crises etc. Quantitative labeling of graph elements have been used in missile guidance codes, radar location codes, coding theory, X-ray crystallography, astronomy, circuit design, communication network etc. Today graph theory is one of the flourishing branches of modern algebra with wide applications to combinatorial problems and to classical algebraic problems.

Graph labeling is one of the fascinating areas of graph theory with wide range of applications. A graph labeling is an assignment of integers to the vertices or the edges or both, subject to certain conditions. If the domain is the

*Correspondence to: D. Praveen Prabhu (Email: prof.praveenprabhu@gmail.com). Department of Mathematics, PPG Institute of Technology, Coimbatore - 641 035, Tamil Nadu, India.

set of vertices we speak about the vertex labeling. If the domain is the set of edges, then the labeling is called edge labeling. Most graph labeling methods trace their origin to the one introduced by Rosa in 1967. Over the past three decades more than 600 papers have been published in graph labeling.

For all other standard terminology and notations, we follow Harary[4]. Graph labeling is used in many areas of sciences and technology. A lot of graph labeling techniques are discussed in [3]. The concept and notation of mean labeling was first introduced by S. Somasundaram and R. Ponraj[10]. S. Somasundaram, S.S. Sandya and T. Pavithra[22] introduced the concept of Lehmer-3 mean labeling of graphs and studied their behavior in [17, 18, 19] and also proved that some Super Lehmer-3 mean labeling and k -Super Lehmer-3 mean labeling graphs [20, 21, 23, 24]. Paramaguru. N and Manirathnam. D[9] proved that some cycle related Lehmer-3 Mean Graphs. S. Somasundaram, R. Ponraj and S.S. Sandhya introduced the concept of Harmonic Mean labeling of graphs in [14] and studied their behavior in [15, 12, 13, 11, 16]. Deepika. R[2] investigated super Lehmer-3 mean labeling of path union related graphs. Meena. S and Mani. R[7] investigated this labeling for some cycle related graphs and they also find some new Lehmer-3 mean graphs in [5, 6]. Meena. S and R. Mani[8] proved root square mean labeling of theta related graphs. Prime cordial labeling for theta graph has been proved by Sugumaran. A and Vishnu Prakash. P[25]. In this paper, we examine about the super Lehmer-3 mean labeling of theta related graphs and some new examples are presented and verified. We now give the definitions which are necessary for the present investigation.

Let $h : V(G) \rightarrow \{1, 2, \dots, p + q\}$ be an injective function. The induced edge labeling $h^*(e = uv)$ is defined by $h^*(e) = \lceil \frac{h(u)^3 + h(v)^3}{h(u)^2 + h(v)^2} \rceil$ or $\lfloor \frac{h(u)^3 + h(v)^3}{h(u)^2 + h(v)^2} \rfloor$, then h is called super Lehmer-3 mean labeling, if $\{h(V(G))\} \cup \{h(e)/e \in E(G)\} = \{1, 2, \dots, p + q\}$. A graph which admits Super Lehmer-3 mean labeling is called Super Lehmer-3 Mean graph. In this paper, we prove that theta related graphs such as θ_τ , $\theta_\tau \odot \overline{K}_1$, $\theta_\tau \odot \overline{K}_2$, $\theta_\tau \odot \overline{K}_3$, $\theta_\tau \odot K_2$, $S(\theta_\tau)$, $(S(\theta_\tau)) \odot \overline{K}_1$, $(S(\theta_\tau)) \odot \overline{K}_2$, $(S(\theta_\tau)) \odot \overline{K}_3$, $(S(\theta_\tau)) \odot K_2$. Subdividing all the path vertices of theta graph such as $\theta_\tau \odot \overline{K}_1$, $\theta_\tau \odot \overline{K}_2$, $\theta_\tau \odot \overline{K}_3$ and $\theta_\tau \odot K_2$ are all super Lehmer-3 mean graphs. Then some special results for harmonic mean labeling of theta graphs also are investigates in this paper.

Definition 1.1

A graph G with p vertices and q edges is called Lehmer-3 mean graph if it is possible to label the vertices $x \in V$ with distinct labels $h(x)$ from $1, 2, \dots, q + 1$ in such a way that when each edge $e = uv$ is labeled with $h(e = uv) = \lceil \frac{h(u)^3 + h(v)^3}{h(u)^2 + h(v)^2} \rceil$ or $\lfloor \frac{h(u)^3 + h(v)^3}{h(u)^2 + h(v)^2} \rfloor$, then the resulting edge labels are distinct. In this case, h is called a **Lehmer-3 mean labeling** of G .

Definition 1.2

Let $h : V(G) \rightarrow \{1, 2, \dots, p + q\}$ be an injective function. The induced edge labeling $h^*(e = uv)$ is defined by $h^*(e) = \lceil \frac{h(u)^3 + h(v)^3}{h(u)^2 + h(v)^2} \rceil$ or $\lfloor \frac{h(u)^3 + h(v)^3}{h(u)^2 + h(v)^2} \rfloor$, then h is called super lehmer-3 mean labeling, if $\{h(V(G))\} \cup \{h(e)/e \in E(G)\} = \{1, 2, \dots, p + q\}$. A graph which admits Super lehmer-3 mean labeling is called **Super Lehmer-3 Mean graph**.

Example 1.2.1 1. Path:



Number of Vertices $p = 6$

Number of Edges $q = 5$

2. Cycle:

Number of Vertices $p = 9$

Number of Edges $q = 9$

Figure 1. Super Lehmer-3 mean labeling of Path P_6 Figure 2. Super Lehmer-3 mean labeling of Cycle C_9 **Definition 1.3**

A graph G with p vertices and q edges is called a **Harmonic mean graph** if it is possible to label the vertices $x \in V$ with distinct labels $f(x)$ from $1, 2, \dots, q+1$ in such a way that when each edge $e = uv$ is labeled with $f(e = uv) = \lceil \frac{2f(u)f(v)}{f(u)+f(v)} \rceil$ or $\lfloor \frac{2f(u)f(v)}{f(u)+f(v)} \rfloor$ then the edge labels are distinct. In this case f is called a **Harmonic mean labeling** of G .

Definition 1.4

A walk in which $x_1x_2 \dots x_\tau$ are distinct is called a **path**. A path on τ vertices is denoted by P_τ .

Definition 1.5

A closed path is called a **cycle**. A cycle on n -vertices is denoted by C_τ .

Definition 1.6

The **Corona of two graphs** G_1 and G_2 is the graph $G = G_1 \odot G_2$ formed by taking one copy of G_1 and $|V(G_1)|$ copies of G_2 where the i^{th} vertex of G_1 is adjacent to every vertex in the i^{th} copy of G_2 .

Definition 1.7

A **subdivision graph** $S(G)$ of a graph G is a graph obtained from G by subdividing all edges of G exactly once.

Example 1.7.1

Path:




Figure 3. G

Subdivision of Path:




Figure 4. $S(G)$

Definition 1.8

A **theta graph** is a block with two non-adjacent vertices of degree 3 and all other vertices of degree 2 is called a theta graph and it is denoted by θ_γ .




Figure 5. Theta Graph θ_τ

Notation

Here $\lceil h \rceil$ is the least integer greater than or equal to h and $\lfloor h \rfloor$ is the greatest integer less than or equal to h .

2. Main Results

In this paper, we investigate the super lehmer-3 mean labeling of theta related graphs.

Theorem 2.1

θ_τ is a super lehmer-3 mean graph.

Proof

Let $G = \theta_\tau$.

Let $x_1x_2 \dots x_\tau$ be the vertices of theta in G .

Let $V(G) = \{x_1x_2 \dots x_\tau\}$

$E(G) = \{x_kx_{k+1}/1 \leq k \leq \tau - 1\} \cup \{x_2x_\tau, x_1x_{\tau-2}\}$.

Define a mapping $h : V(G) \rightarrow \{1, 2, \dots, 2\tau - 1\}$ by

$$h(x_k) = 2k - 1 \text{ for } 1 \leq k \leq \tau.$$

Then the edge labels are

$$h(x_kx_{k+1}) = 2k \text{ for } 1 \leq k \leq \tau - 1$$

$$h(x_1x_{\tau-2}) = \tau + 2$$

$$h(x_2x_\tau) = 2\tau - 1$$

Hence h is a super lehmer-3 mean labeling of G . □

Example 2.1.1

For instance,




Figure 6. Super lehmer-3 mean labeling of θ_7

Theorem 2.2

$S(\theta_\tau)$ is a super lehmer-3 mean graph.

Proof

Let $G = S(\theta_\tau)$.

Let $x_1x_2 \dots x_\tau$ be the vertices of theta in G .

Let $y_1y_2 \dots y_\tau$ be the vertices of subdivision theta graph $S(G)$.

Let $V(G) = \{x_1x_2 \dots x_\tau, y_1y_2 \dots y_\tau\}$

$$E(G) = \{x_kx_{k+1}/1 \leq k \leq \tau\} \cup \{y_kx_k/1 \leq k \leq \tau\} \cup \{y_1x_{\tau-2}, x_2y_{\tau+1}\}$$


Define a mapping $h : V(G) \rightarrow \{1, 2, \dots, 4\tau + 1\}$ by

$$h(x_k) = 4k - 1 \text{ for } 1 \leq k \leq \tau$$

$$h(y_k) = 4k - 3 \text{ for } 1 \leq k \leq \tau$$

$$\begin{aligned} h(x_k y_{k+1}) &= 4k \text{ for } 1 \leq k \leq \tau \\ h(y_k x_k) &= 4k - 2 \text{ for } 1 \leq k \leq \tau \\ h(y_1 x_{\gamma-2}) &= 3\tau - 2 \\ h(x_2 y_{\tau+1}) &= 4\tau - 1 \end{aligned}$$
☐

For instance,



Theorem 2.3

Proof

Let $x_1 x_2 \dots x_\tau$ be the vertices of theta in G .

Let $y_1 y_2 \dots y_\tau$ be the pendant vertices attached at $x_1 x_2 \dots x_\tau$ respectively.

Let $V(G) = \{x_1x_2 \dots x_\tau, y_1y_2 \dots y_\tau\}$

$$E(G) = \{x_k x_{k+1}/1 \leq k \leq \tau - 1\} \cup \{x_k y_k/1 \leq k \leq \tau\} \\ \cup \{x_2 x_\tau, x_1 x_{\tau-2}\}$$

Define a mapping $h : V(G) \rightarrow \{1, 2, \dots, 4\tau - 1\}$ by

$$\begin{aligned} h(x_k) &= 4k - 3 \text{ for } 1 \leq k \leq \tau \\ h(y_k) &= 4k - 1 \text{ for } 1 \leq k \leq \tau \end{aligned}$$

$$\begin{aligned} h(x_k x_{k+1}) &= 4k \text{ for } 1 \leq k \leq \tau - 1 \\ h(x_k y_k) &= 4k - 2 \text{ for } 1 \leq k \leq \tau \\ h(x_1 x_{\tau-2}) &= 2\tau + 3 \\ h(x_2 x_\tau) &= 3\tau - 4 \end{aligned}$$

1

Example 2.3.1

For instance,




Figure 8. Super lehmer-3 mean labeling of $\theta_7 \odot \bar{K}_1$

Theorem 2.4

$(S(\theta_\tau)) \odot \bar{K}_1$ is a super lehmer-3 mean graph.

Proof

Let $G = (S(\theta_\tau)) \odot \bar{K}_1$.

Let $x_1x_2 \dots x_{2\tau+1}$ be the vertices of subdivision theta graph $S(G)$.

Let $y_1y_2 \dots y_{2\tau+1}$ be the pendant vertices attached at $x_1x_2 \dots x_{2\tau+1}$ respectively.

Let $V(G) = \{x_1x_2 \dots x_{2\tau+1}, y_1y_2 \dots y_{2\tau+1}\}$

$$E(G) = \{x_kx_{k+1}/1 \leq k \leq 2\tau\} \cup \{x_ky_k/1 \leq k \leq 2\tau+1\} \\ \cup \{x_1x_{\tau+3}, x_4x_{2\tau+1}\}$$

Define a mapping $h : V(G) \rightarrow \{1, 2, \dots, 8\tau + 3\}$ by

$$h(x_k) = 4k - 3 \text{ for } 1 \leq k \leq 2\tau + 1$$

$$h(y_k) = 4k - 1 \text{ for } 1 \leq k \leq 2\tau + 1$$

Then the edge labels are

$$h(x_kx_{k+1}) = 4k \text{ for } 1 \leq k \leq 2\tau$$

$$h(x_ky_k) = 4k - 2 \text{ for } 1 \leq k \leq 2\tau + 1$$


$$h(x_1x_{\tau+3}) = 5\tau + 3$$

$$h(x_4x_{2\tau+1}) = 8\tau - 1$$

Hence h is a super lehmer-3 mean labeling of G . □

Example 2.4.1

For instance,

Figure 9. Super lehmer-3 mean labeling of $(S(\theta_7)) \odot \bar{K}_1$ **Theorem 2.5**

$\theta_\tau \odot \bar{K}_2$ is a super lehmer-3 mean graph.

Proof

Let $G = \theta_\tau \odot \bar{K}_2$.

Let $x_1 x_2 \dots x_\tau$ be the vertices of theta in G .

Let $y_1 y_2 \dots y_\tau$ and $z_1 z_2 \dots z_\tau$ be the pendant vertices attached at $x_1 x_2 \dots x_\tau$ respectively.

Let $V(G) = \{x_1 x_2 \dots x_\tau, y_1 y_2 \dots y_\tau, z_1 z_2 \dots z_\tau\}$

$$E(G) = \{x_k x_{k+1}/1 \leq k \leq \tau - 1\} \cup \{x_k y_k/1 \leq k \leq \tau\} \\ \cup \{x_k z_k/1 \leq k \leq \tau\} \cup \{x_2 x_\tau, x_1 x_{\tau-2}\}$$

Define a mapping $h : V(G) \rightarrow \{1, 2, \dots, 6\tau - 1\}$ by

$$\begin{aligned} h(x_k) &= 6k - 5 \text{ for } 1 \leq k \leq \tau \\ h(y_k) &= 6k - 3 \text{ for } 1 \leq k \leq \tau \\ h(z_k) &= 6k - 1 \text{ for } 1 \leq k \leq \tau \end{aligned}$$


Then the edge labels are

$$\begin{aligned} h(x_k x_{k+1}) &= 6k \text{ for } 1 \leq k \leq \tau - 4 \\ h(x_k x_{k+1}) &= 6k - 1 \text{ for } \tau - 3 \leq k \leq \tau - 1 \\ h(x_k y_k) &= 6k - 4 \text{ for } 1 \leq k \leq \tau \\ h(x_k z_k) &= 6k - 2 \text{ for } 1 \leq k \leq \tau \\ h(x_1 x_{\tau-2}) &= 3\tau + 4 \\ h(x_2 x_\tau) &= 5\tau + 1 \end{aligned}$$

Hence h is a super lehmer-3 mean labeling of G . □

Example 2.5.1

For instance,

Figure 10. Super lehmer-3 mean labeling of $\theta_7 \odot \bar{K}_2$ **Theorem 2.6**

$(S(\theta_\tau)) \odot \bar{K}_2$ is a super lehmer-3 mean graph.

Proof

Let $G = (S(\theta_\tau)) \odot \bar{K}_2$.

Let $x_1x_2 \dots x_{2\tau+1}$ be the vertices of subdivision theta graph $S(G)$.

Let $y_1y_2 \dots y_{2\tau+1}$ and $z_1z_2 \dots z_{2\tau+1}$ be the pendant vertices attached at $x_1x_2 \dots x_{2\tau+1}$ respectively.

Let $V(G) = \{x_1x_2 \dots x_{2\tau+1}, y_1y_2 \dots y_{2\tau+1}, z_1z_2 \dots z_{2\tau+1}\}$

$$E(G) = \{x_kx_{k+1}/1 \leq k \leq 2\tau\} \cup \{x_ky_k/1 \leq k \leq 2\tau+1\} \\ \cup \{x_kz_k/1 \leq k \leq 2\tau+1\} \cup \{x_1x_{\tau+3}, x_4x_{2\tau+1}\}$$

Define a mapping $h : V(G) \rightarrow \{1, 2, \dots, 13\tau - 2\}$ by

$$\begin{aligned} h(x_k) &= 6k - 5 \text{ for } 1 \leq k \leq 2\tau + 1 \\ h(y_k) &= 6k - 3 \text{ for } 1 \leq k \leq 2\tau + 1 \\ h(z_k) &= 6k - 1 \text{ for } 1 \leq k \leq 2\tau + 1 \end{aligned}$$


Then the edge labels are

$$\begin{aligned} h(x_kx_{k+1}) &= 6k \text{ for } 1 \leq k \leq \tau - 4 \\ h(x_kx_{k+1}) &= 6k - 1 \text{ for } \tau - 3 \leq k \leq 2\tau \\ h(x_ky_k) &= 6k - 4 \text{ for } 1 \leq k \leq 2\tau + 1 \\ h(x_kz_k) &= 6k - 2 \text{ for } 1 \leq k \leq 2\tau + 1 \\ h(x_1x_{\tau+3}) &= 8\tau - 2 \\ h(x_4x_{2\tau+1}) &= 12\tau - 3 \end{aligned}$$

Hence h is a super lehmer-3 mean labeling of G . □

Example 2.6.1

For instance,

Figure 11. Super lehmer-3 mean labeling of $(S(\theta_7)) \odot \bar{K}_2$ **Theorem 2.7**

$\theta_\tau \odot \bar{K}_3$ is a super lehmer-3 mean graph.

Proof

Let $G = \theta_\tau \odot \bar{K}_3$.

Let $x_1 x_2 \dots x_\tau$ be the vertices of theta in G .

Let $y_1 y_2 \dots y_\tau, y'_1 y'_2 \dots y'_\tau$ and $z_1 z_2 \dots z_\tau$ be the pendant vertices attached at $x_1 x_2 \dots x_\tau$ respectively.

Let $V(G) = \{x_1 x_2 \dots x_\tau, y_1 y_2 \dots y_\tau, y'_1 y'_2 \dots y'_\tau, z_1 z_2 \dots z_\tau\}$

$$E(G) = \{x_k x_{k+1} / 1 \leq k \leq \tau - 1\} \cup \{x_k y_k / 1 \leq k \leq \tau\} \\ \cup \{x_k y'_k / 1 \leq k \leq \tau\} \cup \{x_k z_k / 1 \leq k \leq \tau\} \cup \{x_2 x_\tau, x_1 x_{\tau-2}\}$$

Define a mapping $h : V(G) \rightarrow \{1, 2, \dots, 8\tau - 1\}$ by

$$\begin{aligned} h(x_k) &= 8k - 7 \text{ for } 1 \leq k \leq \tau \\ h(y_k) &= 8k - 5 \text{ for } 1 \leq k \leq \tau \\ h(y'_k) &= 8k - 3 \text{ for } 1 \leq k \leq \tau \\ h(z_k) &= 8k - 1 \text{ for } 1 \leq k \leq \tau \end{aligned}$$

Then the edge labels are

$$\begin{aligned} h(x_k x_{k+1}) &= 8k \text{ for } 1 \leq k \leq \tau - 5 \\ h(x_k x_{k+1}) &= 8k - 1 \text{ for } \tau - 4 \leq k \leq \tau - 3 \\ h(x_k x_{k+1}) &= 8k - 2 \text{ for } \tau - 2 \leq k \leq \tau - 1 \\ h(x_k y_k) &= 8k - 6 \text{ for } 1 \leq k \leq \tau \\ h(x_k y'_k) &= 8k - 4 \text{ for } 1 \leq k \leq \tau \\ h(x_k z_k) &= 8k - 2 \text{ for } 1 \leq k \leq \tau - 5 \\ h(x_k z_k) &= 8k - 3 \text{ for } \tau - 4 \leq k \leq \tau \\ h(x_1 x_{\tau-2}) &= 5\tau - 2 \\ h(x_2 x_\tau) &= 7\tau - 2 \end{aligned}$$

☐

For instance,




Figure 12. Super lehmer-3 mean labeling of $\theta_7 \odot \bar{K}_3$

$(S(\theta_\tau)) \odot \bar{K}_3$ is a super lehmer-3 mean graph.

Let $G = (S(\theta_\tau)) \odot \bar{K}_3$.

Let $y_1 y_2 \dots y_{2\tau+1}$, $y'_1 y'_2 \dots y'_{2\tau+1}$ and $z_1 z_2 \dots z_{2\tau+1}$ be the pendant vertices attached at $x_1 x_2 \dots x_{2\tau+1}$ respectively.

Let $V(G) = \{x_1x_2 \dots x_{2\tau+1}, y_1y_2 \dots y_{2\tau+1}, y'_1y'_2 \dots y'_{2\tau+1}, z_1z_2 \dots z_{2\tau+1}$

$$E(G) = \{x_k x_{k+1} / 1 \leq k \leq 2\tau\} \cup \{x_k y_k / 1 \leq k \leq 2\tau + 1\} \\ \cup \{x_k y'_k / 1 \leq k \leq 2\tau + 1\} \cup \{x_k z_k / 1 \leq k \leq 2\tau + 1\} \cup \{x_1 x_{\tau+3}, x_4 x_{2\tau+1}\}$$

Define a mapping $h : V(G) \rightarrow \{1, 2, \dots, 17\tau\}$ by

$$\begin{aligned} h(x_k) &= 8k - 7 \text{ for } 1 \leq k \leq 2\tau + 1 \\ h(y_k) &= 8k - 5 \text{ for } 1 \leq k \leq 2\tau + 1 \\ h(y'_k) &= 8k - 3 \text{ for } 1 \leq k \leq 2\tau + 1 \\ h(z_k) &= 8k - 1 \text{ for } 1 \leq k \leq 2\tau + 1 \end{aligned}$$

Then the edge labels are

$$\begin{aligned}
 h(x_k x_{k+1}) &= 8k \text{ for } 1 \leq k \leq \tau - 5 \\
 h(x_k x_{k+1}) &= 8k - 2 \text{ for } \tau - 4 \leq k \leq 2\tau \\
 h(x_k y_k) &= 8k - 6 \text{ for } 1 \leq k \leq 2\tau + 1 \\
 h(x_k y'_k) &= 8k - 4 \text{ for } 1 \leq k \leq 2\tau - 1 \\
 h(x_k y'_k) &= 8k - 5 \text{ for } 2\tau \leq k \leq 2\tau + 1 \\
 h(x_k z_k) &= 8k - 2 \text{ for } 1 \leq k \leq \tau - 5 \\
 h(x_k z_k) &= 8k - 3 \text{ for } \tau - 4 \leq k \leq 2\tau + 1 \\
 h(x_1 x_{\tau+3}) &= 10\tau + 2 \\
 h(x_4 x_{2\tau+1}) &= 15\tau + 3
 \end{aligned}$$

Hence h is a super lehmer-3 mean labeling of G . □

Example 2.8.1

For instance,




Figure 13. Super lehmer-3 mean labeling of $(S(\theta_7)) \odot \bar{K}_3$

Theorem 2.9

$\theta_\tau \odot K_2$ is a super lehmer-3 mean graph.

Proof

Let $G = \theta_\tau \odot K_2$.

Let $x_1 x_2 \dots x_\tau$ be the vertices of theta in G .

Let $y_1 y_2 \dots y_\tau$ and $z_1 z_2 \dots z_\tau$ be the adjacent vertices attached at $x_1 x_2 \dots x_\tau$ respectively.

Let $V(G) = \{x_1 x_2 \dots x_\tau, y_1 y_2 \dots y_\tau, z_1 z_2 \dots z_\tau\}$

$$\begin{aligned}
 E(G) &= \{x_k x_{k+1} / 1 \leq k \leq \tau - 1\} \cup \{x_k y_k / 1 \leq k \leq \tau\} \\
 &\quad \cup \{x_k z_k / 1 \leq k \leq \tau\} \cup \{y_k z_k / 1 \leq k \leq \tau\} \\
 &\quad \cup \{x_2 x_\tau, x_1 x_{\tau-2}\}
 \end{aligned}$$

Define a mapping $h : V(G) \rightarrow \{1, 2, \dots, 7\tau - 2\}$ by

$$\begin{aligned} h(x_k) &= 6k - 5 \text{ for } 1 \leq k \leq \tau - 3 \\ h(x_k) &= 8k - 13 \text{ for } \tau - 2 \leq k \leq \tau \\ h(y_k) &= 6k - 3 \text{ for } 1 \leq k \leq \tau - 3 \\ h(y_k) &= 8k - 11 \text{ for } \tau - 2 \leq k \leq \tau \\ h(z_k) &= 6k - 1 \text{ for } 1 \leq k \leq \tau - 3 \\ h(z_k) &= 8k - 9 \text{ for } \tau - 2 \leq k \leq \tau \end{aligned}$$

Then the edge labels are

$$\begin{aligned} h(x_k x_{k+1}) &= 6k \text{ for } 1 \leq k \leq \tau - 3 \\ h(x_k x_{k+1}) &= 8k - 8 \text{ for } \tau - 2 \leq k \leq \tau - 1 \\ h(x_k y_k) &= 6k - 4 \text{ for } 1 \leq k \leq \tau - 3 \\ h(x_k y_k) &= 8k - 12 \text{ for } \tau - 2 \leq k \leq \tau \\ h(y_k z_k) &= 6k - 1 \text{ for } 1 \leq k \leq \tau - 3 \\ h(y_k z_k) &= 8k - 9 \text{ for } \tau - 2 \leq k \leq \tau \\ h(x_k z_k) &= 6k - 2 \text{ for } 1 \leq k \leq \tau - 3 \\ h(x_k z_k) &= 8k - 10 \text{ for } \tau - 2 \leq k \leq \tau \\ h(x_1 x_{\tau-2}) &= 4\tau - 2 \\ h(x_2 x_\tau) &= 6\tau \end{aligned}$$

Hence h is a super lehmer-3 mean labeling of G . □

Example 2.9.1

For instance,




Figure 14. Super lehmer-3 mean labeling of $\theta_7 \odot K_2$

Theorem 2.10

$(S(\theta_\tau)) \odot K_2$ is a super lehmer-3 mean graph.

Proof

Let $G = (S(\theta_\tau)) \odot K_2$.

Let $x_1x_2 \dots x_{2\tau+1}$ be the vertices of subdivision theta graph $S(G)$.

Let $y_1y_2 \dots y_{2\tau+1}$ and $z_1z_2 \dots z_{2\tau+1}$ be the adjacent vertices attached at $x_1x_2 \dots x_{2\tau+1}$ respectively.

Let $V(G) = \{x_1x_2 \dots x_{2\tau+1}, y_1y_2 \dots y_{2\tau+1}, z_1z_2 \dots z_{2\tau+1}\}$

$$\begin{aligned} E(G) = & \{x_kx_{k+1}/1 \leq k \leq 2\tau\} \cup \{x_ky_k/1 \leq k \leq 2\tau+1\} \\ & \cup \{y_kz_k/1 \leq k \leq 2\tau+1\} \cup \{x_kz_k/1 \leq k \leq 2\tau+1\} \\ & \cup \{x_1x_{\tau+3}, x_4x_{2\tau+1}\} \end{aligned}$$

Define a mapping $h : V(G) \rightarrow \{1, 2, \dots, 16\tau - 1\}$ by

$$\begin{aligned} h(x_k) &= 6k - 5 \text{ for } 1 \leq k \leq \tau - 3 \\ h(x_k) &= 8k - 13 \text{ for } \tau - 2 \leq k \leq 2\tau + 1 \\ h(y_k) &= 6k - 4 \text{ for } 1 \leq k \leq \tau - 3 \\ h(y_k) &= 8k - 11 \text{ for } \tau - 2 \leq k \leq 2\tau + 1 \\ h(z_k) &= 6k - 1 \text{ for } 1 \leq k \leq \tau - 3 \\ h(z_k) &= 8k - 9 \text{ for } \tau - 2 \leq k \leq 2\tau + 1 \end{aligned}$$

Then the edge labels are

$$\begin{aligned} h(x_kx_{k+1}) &= 6k \text{ for } 1 \leq k \leq \tau - 3 \\ h(x_kx_{k+1}) &= 8k - 8 \text{ for } \tau - 2 \leq k \leq 2\tau \\ h(x_ky_k) &= 6k - 4 \text{ for } 1 \leq k \leq \tau - 3 \\ h(x_ky_k) &= 8k - 12 \text{ for } \tau - 2 \leq k \leq 2\tau + 1 \\ h(y_kz_k) &= 6k - 1 \text{ for } 1 \leq k \leq \tau - 3 \\ h(y_kz_k) &= 8k - 9 \text{ for } \tau - 2 \leq k \leq 2\tau + 1 \\ h(x_kz_k) &= 6k - 2 \text{ for } 1 \leq k \leq \tau - 3 \\ h(x_kz_k) &= 8k - 10 \text{ for } \tau - 2 \leq k \leq 2\tau + 1 \\ h(x_1x_{\tau+3}) &= 10\tau - 4 \\ h(x_4x_{2\tau+1}) &= 15\tau \end{aligned}$$

Hence h is a super lehmer-3 mean labeling of G .

□

Example 2.10.1

For instance,




Figure 15. Super lehmer-3 mean labeling of $(S(\theta_7)) \odot K_2$

Theorem 2.11

The graph G obtained by subdividing all the path vertices of theta graph $\theta_\tau \odot \bar{K}_1$ is a super lehmer - 3 mean graph.

Proof

Let $x_1x_2 \dots x_\tau$ be the vertices of theta in G .

Let $y_1y_2 \dots y_\tau$ be the pendant vertices attached at $x_1x_2 \dots x_\tau$ respectively.

Let $z_1z_2 \dots z_{\tau+1}$ be the vertices obtained by subdividing all the path edges x_kx_{k+1} , $x_1x_{\tau-2}$ and x_2x_τ respectively.

Let $V(G) = \{x_1x_2 \dots x_\tau, y_1y_2 \dots y_\tau, z_1z_2 \dots z_{\tau+1}\}$

$$E(G) = \{z_kx_k/1 \leq k \leq \tau\} \cup \{x_kz_{k+1}/1 \leq k \leq \tau\} \\ \cup \{x_ky_k/1 \leq k \leq \tau\} \cup \{x_2z_{\tau+1}, x_1x_{\tau-2}\}$$

Define a mapping $h : V(G) \rightarrow \{1, 2, \dots, 6\tau + 1\}$ by

$$\begin{aligned} h(x_k) &= 6k - 3 \text{ for } 1 \leq k \leq \tau \\ h(y_k) &= 6k - 1 \text{ for } 1 \leq k \leq \tau \\ h(z_k) &= 6k - 5 \text{ for } 1 \leq k \leq \tau + 1 \end{aligned}$$


Then the edge labels are

$$\begin{aligned} h(z_kx_k) &= 6k - 4 \text{ for } 1 \leq k \leq \tau \\ h(x_kz_{k+1}) &= 6k \text{ for } 1 \leq k \leq \tau \\ h(x_ky_k) &= 6k - 2 \text{ for } 1 \leq k \leq \tau \\ h(z_1x_{\tau-2}) &= 4\tau - 1 \\ h(x_2z_{\tau+1}) &= 6\tau - 1 \end{aligned}$$

Hence h is a super lehmer-3 mean labeling of G . □

Example 2.11.1

For instance,

Figure 16. A Super lehmer-3 mean labeling of Subdividing all the path vertices of theta graph $\theta_7 \odot \bar{K}_1$ *Theorem 2.12*

The graph G obtained by subdividing all the path vertices of theta graph $\theta_\tau \odot \bar{K}_2$ is a super lehmer - 3 mean graph.

Proof

Let $x_1x_2 \dots x_\tau$ be the vertices of theta in G .

Let $y_1y_2 \dots y_\tau$ and $y'_1y'_2 \dots y'_\tau$ be the pendant vertices attached at $x_1x_2 \dots x_\tau$ respectively.

Let $z_1z_2 \dots z_{\tau+1}$ be the vertices obtained by subdividing all the path edges x_kx_{k+1} , $x_1x_{\tau-2}$ and x_2x_τ respectively.

Let $V(G) = \{x_1x_2 \dots x_\tau, y_1y_2 \dots y_\tau, y'_1y'_2 \dots y'_\tau, z_1z_2 \dots z_{\tau+1}\}$

$$\begin{aligned} E(G) = & \{z_kx_k/1 \leq k \leq \tau\} \cup \{x_kz_{k+1}/1 \leq k \leq \tau\} \\ & \cup \{x_ky_k/1 \leq k \leq \tau\} \cup \{x_ky'_k/1 \leq k \leq \tau\} \\ & \cup \{x_2z_{\tau+1}, x_1x_{\tau-2}\} \end{aligned}$$

Define a mapping $h : V(G) \rightarrow \{1, 2, \dots, 8\tau + 1\}$ by

$$\begin{aligned} h(x_k) &= 8k - 5 \text{ for } 1 \leq k \leq \tau \\ h(y_k) &= 8k - 3 \text{ for } 1 \leq k \leq \tau \\ h(y'_k) &= 8k - 1 \text{ for } 1 \leq k \leq \tau \\ h(z_k) &= 8k - 7 \text{ for } 1 \leq k \leq \tau + 1 \end{aligned}$$

Then the edge labels are

$$\begin{aligned} h(z_kx_k) &= 8k - 6 \text{ for } 1 \leq k \leq \tau \\ h(x_kz_{k+1}) &= 8k \text{ for } 1 \leq k \leq \tau - 5 \\ h(x_kz_{k+1}) &= 8k - 7 \text{ for } \tau - 4 \leq k \leq \tau \\ h(x_ky_k) &= 8k - 4 \text{ for } 1 \leq k \leq \tau \\ h(x_ky'_k) &= 8k - 2 \text{ for } 1 \leq k \leq \tau \\ h(z_1x_{\tau-2}) &= 5\tau \\ h(x_2z_{\tau+1}) &= 8\tau \end{aligned}$$

Hence h is a super lehmer-3 mean labeling of G . □

Example 2.12.1

For instance,




Figure 17. A Super lehmer-3 mean labeling of Subdividing all the path vertices of theta graph $\theta_7 \odot \bar{K}_2$

Theorem 2.13

The graph G obtained by subdividing all the path vertices of theta graph $\theta_\tau \odot \bar{K}_3$ is a super lehmer - 3 mean graph.

Proof

Let $x_1x_2 \dots x_\tau$ be the vertices of theta in G .

Let $y_1y_2 \dots y_\tau, y'_1y'_2 \dots y'_\tau$ and $y''_1y''_2 \dots y''_\tau$ be the pendant vertices attached at $x_1x_2 \dots x_\tau$ respectively.

Let $z_1z_2 \dots z_{\tau+1}$ be the vertices obtained by subdividing all the path edges $x_kx_{k+1}, x_1x_{\tau-2}$ and x_2x_τ respectively.

Let $V(G) = \{x_1x_2 \dots x_\tau, y_1y_2 \dots y_\tau, y'_1y'_2 \dots y'_\tau, y''_1y''_2 \dots y''_\tau, z_1z_2 \dots z_{\tau+1}\}$

$$\begin{aligned} E(G) = & \{z_kx_k/1 \leq k \leq \tau\} \cup \{x_kz_{k+1}/1 \leq k \leq \tau\} \\ & \cup \{x_ky_k/1 \leq k \leq \tau\} \cup \{x_ky'_k/1 \leq k \leq \tau\} \\ & \cup \{x_ky''_k/1 \leq k \leq \tau\} \cup \{x_2z_{\tau+1}, x_1x_{\tau-2}\} \end{aligned}$$

Define a mapping $h : V(G) \rightarrow \{1, 2, \dots, 10\tau + 1\}$ by

$$\begin{aligned} h(x_k) &= 10k - 7 \text{ for } 1 \leq k \leq \tau \\ h(y_k) &= 10k - 5 \text{ for } 1 \leq k \leq \tau \\ h(y'_k) &= 10k - 3 \text{ for } 1 \leq k \leq \tau \\ h(y''_k) &= 10k - 1 \text{ for } 1 \leq k \leq \tau \\ h(z_k) &= 10k - 9 \text{ for } 1 \leq k \leq \tau + 1 \end{aligned}$$

Then the edge labels are

$$\begin{aligned}
 h(z_k x_k) &= 10k - 8 \text{ for } 1 \leq k \leq \tau \\
 h(x_1 z_2) &= \tau + 3 \\
 h(x_k z_{k+1}) &= 10k - 2 \text{ for } \tau - 5 \leq k \leq \tau \\
 h(x_k y_k) &= 10k - 6 \text{ for } 1 \leq k \leq \tau \\
 h(x_k y'_k) &= 10k - 4 \text{ for } 1 \leq k \leq \tau \\
 h(x_1 y''_1) &= \tau + 1 \\
 h(x_k y''_k) &= 10k - 3 \text{ for } 2 \leq k \leq \tau \\
 h(z_1 x_{\tau-2}) &= 6\tau + 1 \\
 h(x_2 z_{\tau+1}) &= 10\tau - 1
 \end{aligned}$$

Hence h is a super lehmer-3 mean labeling of G . □

Example 2.13.1

For instance,




Figure 18. A Super lehmer-3 mean labeling of Subdividing all the path vertices of theta graph $\theta_7 \odot \bar{K}_3$

Theorem 2.14

The graph G obtained by subdividing all the path vertices of theta graph $\theta_\tau \odot K_2$ is a super lehmer - 3 mean graph.

Proof

Let $x_1 x_2 \dots x_\tau$ be the vertices of theta in G .

Let $y_1 y_2 \dots y_\tau$ and $y'_1 y'_2 \dots y'_\tau$ be the adjacent vertices attached at $x_1 x_2 \dots x_\tau$ respectively.

Let $z_1 z_2 \dots z_{\tau+1}$ be the vertices obtained by subdividing all the path edges $x_k x_{k+1}$, $x_1 x_{\tau-2}$ and $x_2 x_\tau$ respectively.

Let $V(G) = \{x_1 x_2 \dots x_\tau, y_1 y_2 \dots y_\tau, y'_1 y'_2 \dots y'_\tau, z_1 z_2 \dots z_{\tau+1}\}$

$$\begin{aligned}
 E(G) &= \{z_k x_k / 1 \leq k \leq \tau\} \cup \{x_k z_{k+1} / 1 \leq k \leq \tau\} \\
 &\cup \{x_k y_k / 1 \leq k \leq \tau\} \cup \{x_k y'_k / 1 \leq k \leq \tau\} \\
 &\cup \{y_k y'_k / 1 \leq k \leq \tau\} \cup \{x_2 z_{\tau+1}, x_1 x_{\tau-2}\}
 \end{aligned}$$

Define a mapping $h : V(G) \rightarrow \{1, 2, \dots, 8\tau + 2\}$ by

$$\begin{aligned} h(x_k) &= 8k - 5 \text{ for } 1 \leq k \leq \tau - 4 \\ h(x_k) &= 8k - 4 \text{ for } \tau - 3 \leq k \leq \tau \\ h(y_k) &= 8k - 3 \text{ for } 1 \leq k \leq \tau - 4 \\ h(y_k) &= 8k - 2 \text{ for } \tau - 3 \leq k \leq \tau \\ h(y'_k) &= 8k - 1 \text{ for } 1 \leq k \leq \tau - 4 \\ h(y'_k) &= 8k \text{ for } \tau - 3 \leq k \leq \tau \\ h(z_k) &= 8k - 7 \text{ for } 1 \leq k \leq \tau - 4 \\ h(z_k) &= 8k - 6 \text{ for } \tau - 3 \leq k \leq \tau + 1 \end{aligned}$$

Then the edge labels are

$$\begin{aligned} h(z_k x_k) &= 8k - 6 \text{ for } 1 \leq k \leq \tau - 4 \\ h(z_k x_k) &= 8k - 5 \text{ for } \tau - 3 \leq k \leq \tau \\ h(x_k z_{k+1}) &= 8k \text{ for } 1 \leq k \leq \tau \\ h(x_k y_k) &= 8k - 4 \text{ for } 1 \leq k \leq \tau - 4 \\ h(x_k y_k) &= 8k - 3 \text{ for } \tau - 3 \leq k \leq \tau \\ h(y_k y'_k) &= 8k - 1 \text{ for } 1 \leq k \leq \tau \\ h(x_k y'_k) &= 8k - 2 \text{ for } 1 \leq k \leq \tau \\ h(z_1 x_{\tau-2}) &= 5\tau + 1 \\ h(x_2 z_{\tau+1}) &= 8\tau + 1 \end{aligned}$$

Hence h is a super lehmer-3 mean labeling of G . □

Example 2.14.1

For instance,




Figure 19. A Super lehmer-3 mean labeling of Subdividing all the path vertices of theta graph $\theta_7 \odot K_2$

3. Harmonic Mean Labeling

In this section, we investigate the harmonic mean labeling of theta graphs.

Theorem 3.1

θ_τ is a harmonic mean graph.

Proof

Let $G = \theta_\tau$.

Let $x_1x_2 \dots x_\tau$ be the vertices of theta in G .

Let $V(G) = \{x_1x_2 \dots x_\tau\}$

$E(G) = \{x_kx_{k+1}/1 \leq k \leq \tau - 1\} \cup \{x_2x_\tau, x_1x_{\tau-2}\}$.

Define a mapping $h : V(G) \rightarrow \{1, 2, \dots, \tau + 1\}$ by

$$h(x_k) = k \text{ for } 1 \leq k \leq \tau$$

Then the edge labels are

$$\begin{aligned} h(x_1x_2) &= 1 \\ h(x_kx_{k+1}) &= k + 1 \text{ for } \tau - 5 \leq k \leq \tau - 4 \\ h(x_kx_{k+1}) &= k + 2 \text{ for } \tau - 3 \leq k \leq \tau - 1 \\ h(x_2x_\tau) &= \tau - 2 \\ h(x_1x_{\tau-2}) &= \tau - 5 \end{aligned}$$

Hence h is a super lehmer-3 mean labeling of G . □




Figure 20. Harmonic mean labeling of θ_7

In this future, we investigate the harmonic mean labeling of theta related graphs like as $\theta_\tau \odot \bar{K}_1$, $\theta_\tau \odot \bar{K}_2$, $\theta_\tau \odot \bar{K}_3$, $\theta_\tau \odot K_2$ and etc...

4. Conclusion

It is very interesting to find whether a graph admits super lehmer-3 mean labeling or not. We present fourteen new results on super lehmer-3 mean labeling of theta related graphs. Then one special result investigated in harmonic mean labeling graph. In future, we investigate the graphs like as triangular snake, quadrilateral snake, dragon graph, and etc... The investigation about similar results for various graphs families is an open area of research.

Acknowledgment

The authors wish to thank the anonymous referees for their comments and suggestions.

REFERENCES

1. J. A. Bondy, and U. S. R. Murthy, *Graph theory with Application*, Elsevier Science Publishing Co., New York, 1982.
2. R. Deepika, *Super Lehmer-3 Mean Labeling of Path Union Related Graphs*, Journal of Propulsion Technology, vol. 44, no. 02, pp. 1425–1430, 2023.
3. J. A. Gallian, *A Dynamic Survey of Graph Labeling*, The Electronic Journal of Combinatorics, 17#DS6, 2010.
4. F. Harary, *Graph Theory*, Narosa Publishing House Reading, New Delhi, 1988.
5. C. Jaisankar, R. Mani, and M. Manivel, *Lehmer-3 Mean Labeling of Some New Graphs*, Malaya Journal of Matematik, vol. S, no. 1, pp. 188–190, 2021.
6. C. Jaisankar, M. Manivel, and E. Keerthana, *Lehmer-3 mean labeling of dragon graphs*, Malaya Journal of Matematik, vol. S, no. 1, pp. 290–293, 2021.
7. S. Meena, and R. Mani, *Root Square Mean labeling of Some Cycle Related Graphs*, International Journal for Science and Advance Research Technology, vol. 5, no. 7, pp. 786–789, 2019.
8. S. Meena, and R. Mani, *Root Square Mean labeling of Theta Related Graphs*, Mathematical Statistician and Engineering Applications, vol. 71, no. 4, pp. 13049–13070, 2022.
9. N. Paramaguru, and D. Manirathnam, *Lehmer-3 Mean Labeling of Bicyclic Graphs*, Advances in Mechanics, vol. 12, no. 1, pp. 274–286, 2024.
10. R. Ponraj, and S. Somasundaram, *Mean Labeling of Graphs*, National Academy of Science Letters, vol. 26, pp. 210–213, 2003.
11. S. S. Sandhya, C. Jayasekaran, and C. S. David Raj, *Some New Families of Harmonic Mean Graphs*, International Journal of Mathematica Research, vol. 5, no. 1, pp. 223–232, 2013.
12. S. S. Sandhya, C. Jayasekaran, and C. D. Raj, *Harmonic Mean Labeling of Degree Splitting Graphs*, Bulletin of Pure and Applied Science, vol. 32E, pp. 99–112, 2013.
13. S. S. Sandhya, S. Somasundaram, and R. Ponraj, *Some More Results on Harmonic Mean Graphs*, Journal of Mathematics Research, vol. 4, pp. 21–29, 2004.
14. S. S. Sandhya, S. Somasundaram, and R. Ponraj, *Some Results on Harmonic Mean Graphs*, International Journal Contemporary Mathematical Sciences, vol. 7, no. 4, pp. 197–208, 2012.
15. S. S. Sandhya, S. Somasundaram, and R. Ponraj, *Some More Results on Harmonic Mean Graphs*, Journal of Mathematics Research, vol. 4, no. 1, pp. 21–29, 2012.
16. S. S. Sandhya, Somasundaram, and J. Rajeshini Golda, *Subdivision of Contra Harmonic Mean Graphs*, International Journal of Mathematics and its Applications, vol. 5, no. 2-c pp. 391–401, 2017.
17. S. Somasundaram, S.S. Sandhya, and T. S. Pavithra, *Some More Results on Lehmer-3 Mean Labeling of Graphs*, Global Journal of Theoretical and Applied Mathematics Science, vol. 6, no. 1, 67–77.
18. S. Somasundaram, S.S. Sandhya, and T. S. Pavithra, *Lehmer-3 Mean Labeling of Some New Disconnected Graphs*, International Journal of Mathematics Trends and Technology (IJMTT), vol. 35, no. 1, 1–14, July 2016.
19. S. Somasundaram, S.S. Sandhya, and T. S. Pavithra, *Lehmer-3 Mean Labeling of Some Disconnected Graphs*, International Journal of Mathematics Research, vol. 8, no. 2, 133–142, 2016.
20. S. Somasundaram, S.S. Sandhya, and T. S. Pavithra, *Super Lehmer-3 Mean Labeling*, Journal of Mathematics Research, vol. 8, no. 5, 29–36, 2016.
21. S. Somasundaram, S.S. Sandhya, and T. S. Pavithra, *Super Lehmer-3 Mean Labeling*, Journal of Mathematics Research, vol.8, no. 5, 29–36, October 2016.
22. S. Somasundaram, S.S. Sandhya, and T. S. Pavithra, *Lehmer-3 Mean Labeling of Graphs*, International Journal of Mathematical Forum, vol. 12, no. 17, 819–825, 2017.
23. S. Somasundaram, S.S. Sandhya, and T. S. Pavithra, *Super Lehmer-3 Mean Number of Graphs*, Global Journal of Pure and Applied Mathematics, vol. 13, no. 10, 7377–7385, 2017.
24. S. Somasundaram, S.S. Sandhya, and T. S. Pavithra, *k-Super Lehmer-3 Mean Graph*, International Journal of Scientific and Innovative Mathematical Research (IJSIMR), vol. 5, no. 5, 1–7, 2017.
25. A. Sugumaran, and P. Vishnu Prakash, *Prime Cordial Labeling of Theta Graphs*, Annals of Pure and Applied Mathematics, vol. 14, no. 3, 379–386, 2017.