

Bipolar Valued Vague Subfields of a Field by Mapping

K. Bala Bavithra^{1,*}, M. Muthusamy², K. Arjunan³

¹ *Department of Mathematics, Sonameenal Arts and Science College (Affiliated to Alagappa University, Karaikudi), Mudukulathur – 623704, Tamil Nadu, India. Email: kpavi94pk@gmail.com*

² *Department of Mathematics, Dr. Zakir Husain College (Affiliated to Alagappa University, Karaikudi), Ilayangudi – 630702, Tamil Nadu, India. Email: msamy0207@yahoo.com*

³ *Department of Mathematics, Alagappa Govt. Arts College (Affiliated to Alagappa University, Karaikudi), Karaikudi – 630003, Tamil Nadu, India. Email: arjunan.karmegam@gmail.com*

Abstract The fuzzy concept dives from the crisp, so a niche obtained in the world. A substantial achievement was gotten in fuzzy from many research people. In this way, we are trying to work in the extension of fuzzy with help of function. Here bipolar valued vague subfield of a field is introduced and defined by the previous work; a crucial part of bipolar valued vague subfield of a field is explored that homomorphism and translations are used in this field and these two are vital role in algebra. In this paper, statements and their proof are given as detail.

Keywords Fuzzy set, vague set, bipolar valued vague set, bipolar valued vague subfield, pseudo bipolar valued vague coset, translation.

AMS 2010 subject classifications 03F55, 06D27, 08A72, 16Y60

DOI: 10.19139/soic-2310-5070-2972

Introduction

The crisp set was generalized into many valued logic that was fuzzy set, it was introduced by L.A.Zadeh in 1965. Succeeding years, fuzzy sets grew in different ways. The following are extensions of fuzzy sets: vague set, intuitionistic fuzzy set, bipolar valued fuzzy set, etc. Rosenfeld [15], bipolar valued fuzzy subset by Zhang [19], vague group by Biswas [5], bipolar vague set by Cicily Flora and Arockiarani [6], bipolar valued fuzzy subgroup by Anitha et al. [2], bipolar valued vague subfield by Bala Bavithra et al. [3], and works by Deepa et al. [7, 8, 9] introduced bipolar valued vague subrings of a ring and their properties. In a similar way, other works [1, 4, 12, 13, 14, 16, 17, 18] were useful to write this paper. It is necessary to introduce this paper in the current growth of human life. Upto 16th century we are working in two valued logic, after the introduction of fuzzy set, the growth of people life has been being grown with respect to the many valued logic. We are giving some examples, theorems and characterization of BVVSF in this paper.

1. Preliminaries

Definition 1.1. [18] A map $R : G \rightarrow [0, 1]$ is called a fuzzy subset of G .

*Correspondence to: K. Bala Bavithra (E-mail: kpavi94pk@gmail.com). Department of Mathematics, Sonameenal Arts and Science College (Affiliated to Alagappa University, Karaikudi), Mudukulathur – 623704, Tamil Nadu, India.

Definition 1.2. [10] The ordered structure $\mathcal{U} = \{(z, [\mathcal{U}_T(z), 1 - \mathcal{U}_F(z)]) : z \in W\}$ is called a vague set of W , where $\mathcal{U}_T : W \rightarrow [0, 1]$ is a truth membership map and $\mathcal{U}_F : W \rightarrow [0, 1]$ is a false membership map, such that $\mathcal{U}_T(z) + \mathcal{U}_F(z) \leq 1$, for all z in W .

Definition 1.3. [10] The interval $[\mathcal{U}_T(z), 1 - \mathcal{U}_F(z)]$ is called the vague value of z in \mathcal{U} and it is denoted by $\mathcal{U}(z)$, i.e., $\mathcal{U}(z) = [\mathcal{U}_T(z), 1 - \mathcal{U}_F(z)]$.

Example 1.4. Let $\mathcal{U} = \{< z, [0.04, 0.07] >, < v, [0.02, 0.06] >, < n, [0.03, 0.08] >\}$ is a vague set of $R = \{z, v, n\}$.

Definition 1.5. [19] The ordered structure $T = \{(z, T^+(z), T^-(z)) : z \in W\}$ is called a bipolar valued fuzzy subset of W , where $T^+ : W \rightarrow [0, 1]$ is a positive membership map and $T^- : W \rightarrow [-1, 0]$ is a negative membership map.

Definition 1.6. [6] The ordered structure $\mathcal{U} = \{(z, [\mathcal{U}_T^+(z), 1 - \mathcal{U}_F^+(z)], [-1 - \mathcal{U}_T^-(z), \mathcal{U}_F^-(z)]) : z \in W\}$ is called a bipolar valued vague subset of W , where $\mathcal{U}_T^+ : W \rightarrow [0, 1]$, $\mathcal{U}_F^+ : W \rightarrow [0, 1]$, $\mathcal{U}_T^- : W \rightarrow [-1, 0]$, and $\mathcal{U}_F^- : W \rightarrow [-1, 0]$ are mapping such that $\mathcal{U}_T^+(z) + \mathcal{U}_F^+(z) \leq 1$, $-1 \leq \mathcal{U}_T^-(z) + \mathcal{U}_F^-(z)$, for all z in W . Bipolar valued vague subset \mathcal{U} is denoted as $\mathcal{U} = \{(z, [\mathcal{U}^+(z), \mathcal{U}^-(z)]) : z \in W\}$, where $\mathcal{U}^+(z) = [\mathcal{U}_T^+(z), 1 - \mathcal{U}_F^+(z)]$ and $\mathcal{U}^-(z) = [-1 - \mathcal{U}_T^-(z), \mathcal{U}_F^-(z)]$. It is denoted as $B_{VV}S$.

Example 1.7. Let $\mathcal{U} = \{< z, [0.05, 0.07], [-0.05, -0.02] >, < v, [0.04, 0.08], [-0.06, -0.03] >, < n, [0.14, 0.19], [-0.25, -0.22] >\}$ is a bipolar value vague set of $R = \{z, v, n\}$.

Definition 1.8. [6] Let $\mathcal{U} = (\mathcal{U}^+, \mathcal{U}^-)$ and $G = (G^+, G^-)$ be two bipolar valued vague sets on W . Then

1. $\mathcal{U} \subset G$ if and only if $\mathcal{U}^+(z) \leq G^+(z)$ and $\mathcal{U}^-(z) \geq G^-(z)$, for all $z \in W$.
2. $\mathcal{U} \cap G = \{< z, \text{rmin}(\mathcal{U}^+(z), G^+(z)), \text{rmax}(\mathcal{U}^-(z), G^-(z)) > / z \in W\}$.

Definition 1.9. [3] A $B_{VV}S$ $C = \langle C^+, C^- \rangle$ of a field K is said to be a bipolar valued vague subfield C of $K(B_{VV}SF)$ if C has,

1. $C^+(y - w) \geq \text{rmin}\{C^+(y), C^+(w)\}$,
2. $C^+(yw) \geq \text{rmin}\{C^+(y), C^+(w)\}$,
3. $C^-(y - w) \leq \text{rmax}\{C^-(y), C^-(w)\}$,
4. $C^-(yw) \leq \text{rmax}\{C^-(y), C^-(w)\}$ for all $y, w \in K$,
5. $C^+(y^{-1}) \geq C^+(y) \forall y \neq 0 \in K$,
6. $C^-(y^{-1}) \leq C^-(y) \forall y \neq 0 \in K$,

where 0 is an first operation identity element of K , $\text{rmin}\{[r, s], [t, u]\} = [\min\{r, t\}, \min\{s, u\}]$ and $\text{rmax}\{[r, s], [t, u]\} = [\max\{r, t\}, \max\{s, u\}]$.

Example 1.10. $C^+(y)=[0.36, 0.38]$ for $y \in i2i$, $C^-(y)=[-0.39, -0.36]$ for $y \in i2i$, $C^+(y)=[0.35, 0.37]$, $C^-(y)=[-0.38, -0.35]$ for $y \in R-i2i$ is a $B_{VV}SF$ of the field R .

Definition 1.11. [6] Let $A = \langle A^+, A^- \rangle$ and $W = \langle W^+, W^- \rangle$ be $B_{VV}S$ s of the sets V_1 and V_2 respectively. The product of A and W , denoted by $A \times W$, is defined as $A \times W = \{< (\chi, \zeta), (A \times W)^+(\chi, \zeta), (A \times W)^-(\chi, \zeta) > / \forall (\chi, \zeta) \in V_1 \times V_2\}$, where $(A \times W)^+(\chi, \zeta) = \text{rmin}\{A^+(\chi), W^+(\zeta)\}$ and $(A \times W)^-(\chi, \zeta) = \text{rmax}\{A^-(\chi), W^-(\zeta)\}$.

Definition 1.12. [8] Let X and Y be any two nonempty sets and $f : X \rightarrow Y$ be a mapping. If $A = \langle V_A^+, V_A^- \rangle$ be a bipolar valued vague set in X , then the image of A under f , denoted by $f(A) = B = \langle V_B^+, V_B^- \rangle$ (say), is $B_{VV}S$ in Y defined by $V_B^+(y) = \text{rsup}_{x \in f^{-1}(y)} V_A^+(x)$, $V_B^-(y) = \text{rinf}_{x \in f^{-1}(y)} V_A^-(x)$ if $f^{-1}(y) \neq \phi$ for all $y \in Y$ and $V_B^+(y) = [0, 0]$, $V_B^-(y) = [-1, -1]$ otherwise. If $B = \langle V_B^+, V_B^- \rangle$ is a $B_{VV}S$ in Y , then the preimage of B under f , denoted by $f^{-1}(B) = A = \langle V_A^+, V_A^- \rangle$ (say), is the $B_{VV}S$ in X defined by $V_A^+(x) = V_B^+(f(x))$, $V_A^-(x) = V_B^-(f(x))$ for all $x \in X$.

Definition 1.13. [7] Let $C = \langle V_C^+, V_C^- \rangle$ be a $B_{VV}SF$ of a field F_1 and $s \in F_1$. Then the pseudo bipolar valued vague coset $(sC)^p = \langle (sV_C^+)^{V_p^+}, (sV_C^-)^{V_p^-} \rangle$ is defined by $(sV_C^+)^{V_p^+}(a) = V_p^+(s)V_C^+(a)$ and $(sV_C^-)^{V_p^-}(a) = -V_p^-(s)V_C^-(a)$ for every $a \in F_1$ and $p \in P$, where P is a collection of $B_{VV}SSs$ of F_1 .

Definition 1.14. [9] Let $A = \langle V_A^+, V_A^- \rangle$ be a $B_{VV}S$ of a set X . Then the height $H(A) = \langle H(V_A^+), H(V_A^-) \rangle$ is defined as $H(V_A^+) = \text{rsup}V_A^+(x)$ and $H(V_A^-) = \text{rinf}V_A^-(x)$ for all $x \in X$.

Definition 1.15. [9] Let $A = \langle V_A^+, V_A^- \rangle$ be a $B_{VV}S$ of a set X . Then A is called bipolar valued normal vague subset of X if $H(V_A^+) = [1]$ and $H(V_A^-) = [-1]$.

Definition 1.16. [9] Let $A = \langle V_A^+, V_A^- \rangle$ be a $B_{VV}S$ of X . Then ${}^\circ A = \langle {}^\circ V_A^+, {}^\circ V_A^- \rangle$ is defined as ${}^\circ V_A^+(x) = V_A^+(x)H(V_A^+)$ and ${}^\circ V_A^-(x) = -V_A^-(x)H(V_A^-)$ for all x in X .

Definition 1.17. [9] Let $A = \langle V_A^+, V_A^- \rangle$ be a $B_{VV}S$ of X . Then ${}^\Delta A = \langle {}^\Delta V_A^+, {}^\Delta V_A^- \rangle$ is defined as ${}^\Delta V_A^+(x) = V_A^+(x)/H(V_A^+)$ and ${}^\Delta V_A^-(x) = -V_A^-(x)/H(V_A^-)$ for all x in X .

2. Bipolar Valued Vague Subfield of a Field

Theorem 2.1

If $N = \langle N^+, N^- \rangle$ and $V = \langle V^+, V^- \rangle$ are $B_{VV}SFs$ of fields F_1 and F_2 , respectively, then $N \times V$ is a $B_{VV}SF$ of the field of $F_1 \times F_2$.

Proof

Let $\rho, d \in F_1$ and $\zeta, \xi \in F_2$. Then $(\rho, \zeta), (d, \xi) \in F_1 \times F_2$. Then

$$\begin{aligned} (N \times V)^+[(\rho, \zeta) - (d, \xi)] &= (N \times V)^+[(\rho - d, \zeta - \xi)] \\ &= \text{rmin}\{N^+(\rho - d), V^+(\zeta - \xi)\} \\ &\geq \text{rmin}\{\text{rmin}\{N^+(\rho), N^+(d)\}, \text{rmin}\{V^+(\zeta), V^+(\xi)\}\} \\ &= \text{rmin}\{\text{rmin}\{N^+(\rho), V^+(\zeta)\}, \text{rmin}\{N^+(d), V^+(\xi)\}\} \\ &= \text{rmin}\{(N \times V)^+(\rho, \zeta), (N \times V)^+(d, \xi)\} \end{aligned}$$

for all $(\rho, \zeta), (d, \xi) \in F_1 \times F_2$. And

$$\begin{aligned} (N \times V)^+[(\rho, \zeta)(d, \xi)^{-1}] &= (N \times V)^+[(\rho d^{-1}, \zeta \xi^{-1})] \\ &= \text{rmin}\{N^+(\rho d^{-1}), V^+(\zeta \xi^{-1})\} \\ &\geq \text{rmin}\{\text{rmin}\{N^+(\rho), N^+(d)\}, \text{rmin}\{V^+(\zeta), V^+(\xi)\}\} \\ &= \text{rmin}\{\text{rmin}\{N^+(\rho), V^+(\zeta)\}, \text{rmin}\{N^+(d), V^+(\xi)\}\} \\ &= \text{rmin}\{(N \times V)^+(\rho, \zeta), (N \times V)^+(d, \xi)\} \end{aligned}$$

for all $(\rho, \zeta), (d, \xi) \in F_1 \times F_2$. Also

$$\begin{aligned} (N \times V)^-[(\rho, \zeta) - (d, \xi)] &= (N \times V)^-[(\rho - d, \zeta - \xi)] \\ &= \text{rmax}\{N^-(\rho - d), V^-(\zeta - \xi)\} \\ &\leq \text{rmax}\{\text{rmax}\{N^-(\rho), N^-(d)\}, \text{rmax}\{V^-(\zeta), V^-(\xi)\}\} \\ &= \text{rmax}\{\text{rmax}\{N^-(\rho), V^-(\zeta)\}, \text{rmax}\{N^-(d), V^-(\xi)\}\} \\ &= \text{rmax}\{(N \times V)^-(\rho, \zeta), (N \times V)^-(d, \xi)\} \end{aligned}$$

for all $(\rho, \zeta), (d, \xi) \in F_1 \times F_2$. And

$$\begin{aligned}
 (N \times V)^-[(\rho, \zeta)(d, \xi)^{-1}] &= (N \times V)^-[(\rho d^{-1}, \zeta \xi^{-1})] \\
 &= \text{rmax}\{N^-(\rho d^{-1}), V^-(\zeta \xi^{-1})\} \\
 &\leq \text{rmax}\{\text{rmax}\{N^-(\rho), N^-(d)\}, \text{rmax}\{V^-(\zeta), V^-(\xi)\}\} \\
 &= \text{rmax}\{r \max\{N^-(\rho), V^-(\zeta)\}, \text{rmax}\{N^-(d), V^-(\xi)\}\} \\
 &= \text{rmax}\{(N \times V)^-(\rho, \zeta), (N \times V)^-(d, \xi)\}
 \end{aligned}$$

for all $(\rho, \zeta), (d, \xi) \in F_1 \times F_2$. Hence $N \times V$ is a B_{VV} SFs of $F_1 \times F_2$. \square

Theorem 2.2

Let $C = (V_C^+, V_C^-)$ be a B_{VV} SF of a field F_1 . Then the pseudo bipolar valued vague coset $(sC)^p$ is a B_{VV} SF of F_1 , for every $s \in F_1$ and $p \in P$, where P is a collection of B_{VV} SSs of F_1 .

Proof

Let $a, b \in F_1$,

$$\begin{aligned}
 (V_{sC}^+)^{V_p^+}(a - b) &= V_p^+(s)V_C^+(a - b) \\
 &\geq V_p^+(s) \text{rmin}\{V_C^+(a), V_C^+(b)\} \\
 &= \text{rmin}\{V_p^+(s)V_C^+(a), V_p^+(s)V_C^+(b)\} \\
 &= \text{rmin}\{(V_{sC}^+)^{V_p^+}(a), (V_{sC}^+)^{V_p^+}(b)\}
 \end{aligned}$$

for all $a, b \in F_1$. And

$$\begin{aligned}
 (V_{sC}^+)^{V_p^+}(ab^{-1}) &= V_p^+(s)V_C^+(ab^{-1}) \\
 &\geq V_p^+(s) \text{rmin}\{V_C^+(a), V_C^+(b)\} \\
 &= \text{rmin}\{V_p^+(s)V_C^+(a), V_p^+(s)V_C^+(b)\} \\
 &= \text{rmin}\{(V_{sC}^+)^{V_p^+}(a), (V_{sC}^+)^{V_p^+}(b)\}
 \end{aligned}$$

for all $a, b \in F_1$. Also

$$\begin{aligned}
 (V_{sC}^-)^{V_p^-}(a - b) &= V_p^-(s)V_C^-(a - b) \\
 &\leq V_p^-(s) \text{rmax}\{V_C^-(a), V_C^-(b)\} \\
 &= \text{rmax}\{V_p^-(s)V_C^-(a), V_p^-(s)V_C^-(b)\} \\
 &= \text{rmax}\{(V_{sC}^-)^{V_p^-}(a), (V_{sC}^-)^{V_p^-}(b)\}
 \end{aligned}$$

for all $a, b \in F_1$. And

$$\begin{aligned}
 (V_{sC}^-)^{V_p^-}(ab^{-1}) &= V_p^-(s)V_C^-(ab^{-1}) \\
 &\leq V_p^-(s) \text{rmax}\{V_C^-(a), V_C^-(b)\} \\
 &= \text{rmax}\{V_p^-(s)V_C^-(a), V_p^-(s)V_C^-(b)\} \\
 &= \text{rmax}\{(V_{sC}^-)^{V_p^-}(a), (V_{sC}^-)^{V_p^-}(b)\}
 \end{aligned}$$

for all $a, b \in F_1$. Hence $(sC)^p$ is a B_{VV} SF of F_1 . \square

Theorem 2.3

Let C be a B_{VV} SS of a field F_1 . Then C is a B_{VV} SF of F_1 if and only if each (V_C^+, V_C^-) is a B_V -fuzzy subfield of F_1 .

Proof

Let $a, b \in F_1$. Suppose C is a $B_{VV}SF$ of F_1 ,

$$\begin{aligned} V_C^+(a-b) &\geq \text{rmin}\{V_C^+(a), V_C^+(b)\} \\ V_C^+(ab^{-1}) &\geq \text{rmin}\{V_C^+(a), V_C^+(b)\} \\ V_C^-(a-b) &\leq \text{rmax}\{V_C^-(a), V_C^-(b)\} \\ V_C^-(ab^{-1}) &\leq \text{rmax}\{V_C^-(a), V_C^-(b)\} \end{aligned}$$

Hence each (V_C^+, V_C^-) is a B_V -fuzzy subfield of F_1 .

Conversely, assume that each (V_C^+, V_C^-) is a B_V -fuzzy subfield of F_1 . As per the definition of $B_{VV}SF$ of F_1 , C is a $B_{VV}SF$ of F_1 . \square

Theorem 2.4

If $K = \langle V_K^+, V_K^- \rangle$ is a $B_{VV}SF$ of a field F_1 , then ${}^\circ K = \langle {}^\circ V_K^+, {}^\circ V_K^- \rangle$ is a $B_{VV}SF$ of F_1 .

Proof

For any $u, v \in F_1$,

$$\begin{aligned} {}^\circ V_k^+(u-v) &= V_k^+(u-v)H(V_k^+) \\ &\geq \text{rmin}\{V_k^+(u), V_k^+(v)\}H(V_k^+) \\ &= \text{rmin}\{V_k^+(u)H(V_k^+), V_k^+(v)H(V_k^+)\} \\ &= \text{rmin}\{{}^\circ V_k^+(u), {}^\circ V_k^+(v)\} \end{aligned}$$

$\forall u, v \in F_1$.

And

$$\begin{aligned} {}^\circ V_k^+(uv^{-1}) &= V_k^+(uv^{-1})H(V_k^+) \\ &\geq \text{rmin}\{V_k^+(u), V_k^+(v)\}H(V_k^+) \\ &= \text{rmin}\{V_k^+(u)H(V_k^+), V_k^+(v)H(V_k^+)\} \\ &= \text{rmin}\{{}^\circ V_k^+(u), {}^\circ V_k^+(v)\} \end{aligned}$$

$\forall u, v \in F_1$. Also

$$\begin{aligned} {}^\circ V_k^-(u-v) &= -V_k^-(u-v)H(V_k^-) \\ &\leq -\text{rmax}\{V_k^-(u), V_k^-(v)\}H(V_k^-) \\ &= \text{rmax}\{-V_k^-(u)H(V_k^-), -V_k^-(v)H(V_k^-)\} \\ &= \text{rmax}\{{}^\circ V_k^-(u), {}^\circ V_k^-(v)\} \end{aligned}$$

$\forall u, v \in F_1$. And

$$\begin{aligned} {}^\circ V_k^-(uv^{-1}) &= -V_k^-(uv^{-1})H(V_k^-) \\ &\leq -\text{rmax}\{V_k^-(u), V_k^-(v)\}H(V_k^-) \\ &= \text{rmax}\{-V_k^-(u)H(V_k^-), -V_k^-(v)H(V_k^-)\} \\ &= \text{rmax}\{{}^\circ V_k^-(u), {}^\circ V_k^-(v)\} \end{aligned}$$

$\forall u, v \in F_1$. Hence ${}^\circ K$ is a $B_{VV}SF$ of F_1 . \square

Theorem 2.5

If $K = \langle V_K^+, V_K^- \rangle$ is a $B_{VV}SF$ of a field F_1 , then:

- (i) if $H(V_K^+) < [1]$, then ${}^\circ V_K^+ < V_K^+$;
- (ii) if $H(V_K^-) > [-1]$, then ${}^\circ V_K^- > V_K^-$;
- (iii) if $H(V_K^+) < [1]$ and $H(V_K^-) > [-1]$, then ${}^\circ K < K$.

Proof

(i), (ii) and (iii) are trivial. □

Theorem 2.6

If $K = \langle V_K^+, V_K^- \rangle$ is a $\mathbf{B}_{VV}SF$ of a field F_1 , then ${}^\Delta K = \langle {}^\Delta V_K^+, {}^\Delta V_K^- \rangle$ is a $\mathbf{B}_{VV}SF$ of F_1 .

Proof

For any $u, v \in F_1$, Then

$$\begin{aligned} {}^\Delta V_K^+(u-v) &= V_K^+(u-v)/H(V_K^+) \\ &\geq \text{rmin}\{V_K^+(u), V_K^+(v)\}/H(V_K^+) \\ &= \text{rmin}\{V_K^+(u)/H(V_K^+), V_K^+(v)/H(V_K^+)\} \\ &= \text{rmin}\{{}^\Delta V_K^+(u), {}^\Delta V_K^+(v)\} \end{aligned}$$

$\forall u, v \in F_1$.

And

$$\begin{aligned} {}^\Delta V_K^+(uv^{-1}) &= V_K^+(uv^{-1})/H(V_K^+) \\ &\geq \text{rmin}\{V_K^+(u), V_K^+(v)\}/H(V_K^+) \\ &= \text{rmin}\{V_K^+(u)/H(V_K^+), V_K^+(v)/H(V_K^+)\} \\ &= \text{rmin}\{{}^\Delta V_K^+(u), {}^\Delta V_K^+(v)\} \end{aligned}$$

$\forall u, v \in F_1$. Also

$$\begin{aligned} {}^\Delta V_K^-(u-v) &= -V_K^-(u-v)/H(V_K^-) \\ &\leq -\text{rmax}\{V_K^-(u), V_K^-(v)\}/H(V_K^-) \\ &= \text{rmax}\{-V_K^-(u)/H(V_K^-), -V_K^-(v)/H(V_K^-)\} \\ &= \text{rmax}\{{}^\Delta V_K^-(u), {}^\Delta V_K^-(v)\} \end{aligned}$$

$\forall u, v \in F_1$. And

$$\begin{aligned} {}^\Delta V_K^-(uv^{-1}) &= -V_K^-(uv^{-1})/H(V_K^-) \\ &\leq -\text{rmax}\{V_K^-(u), V_K^-(v)\}/H(V_K^-) \\ &= \text{rmax}\{-V_K^-(u)/H(V_K^-), -V_K^-(v)/H(V_K^-)\} \\ &= \text{rmax}\{{}^\Delta V_K^-(u), {}^\Delta V_K^-(v)\} \end{aligned}$$

$\forall u, v \in F_1$. Hence ${}^\Delta K$ is a $\mathbf{B}_{VV}SF$ of F_1 . □

Theorem 2.7

If $K = \langle V_K^+, V_K^- \rangle$ is a $\mathbf{B}_{VV}SF$ of a field F_1 , then:

- (i) If $H(V_K^+) < [1]$, then ${}^\Delta V_K^+ > V_K^+$;
- (ii) If $H(V_K^-) > [-1]$, then ${}^\Delta V_K^- < V_K^-$;
- (iii) If $H(V_K^+) < [1]$ and $H(V_K^-) > [-1]$, then ${}^\Delta K > K$;
- (iv) If $H(V_K^+) < [1]$ and $H(V_K^-) > [-1]$, then ${}^\Delta K$ is a normal $\mathbf{B}_{VV}SF$ of F_1 .

Proof

(i), (ii), (iii), and (iv) are trivial. □

Theorem 2.8

If $K = \langle V_K^+, V_K^- \rangle$ is a normal $\mathbf{B}_{VV}SF$ of a field F_1 , then:

- (i) $\circ K = K$,
- (ii) $\Delta K = K$.

Proof

It can be easily proved. □

3. Homomorphism on Bipolar Valued Vague Subfield of a Field

Theorem 3.1

The homomorphic image of a $\mathbf{B}_{VV}SF$ of a field F_1 is a $\mathbf{B}_{VV}SF$ of a field F_2 .

Proof

Let $f : F_1 \rightarrow F_2$ be a homomorphism. Let $V = f(A) = \langle V_V^+, V_V^- \rangle$, where $A = \langle V_A^+, V_A^- \rangle$ is a $\mathbf{B}_{VV}SF$ of F_1 . We have to prove that V is a $\mathbf{B}_{VV}SF$ of F_2 . Now for $f(x), f(y) \in F_2$,

$$V_V^+(f(x) - f(y)) = V_V^+(f(x - y)) \geq V_A^+(x - y) \geq \text{rmin}\{V_A^+(x), V_A^+(y)\} = \text{rmin}\{V_V^+(f(x)), V_V^+(f(y))\},$$

which implies that

$$V_V^+(f(x) - f(y)) \geq \text{rmin}\{V_V^+(f(x)), V_V^+(f(y))\}.$$

Also,

$$V_V^+(f(x)f(y)^{-1}) = V_V^+(f(xy^{-1})) \geq V_A^+(xy^{-1}) \geq \text{rmin}\{V_A^+(x), V_A^+(y)\} = \text{rmin}\{V_V^+(f(x)), V_V^+(f(y))\},$$

which implies that

$$V_V^+(f(x)f(y)^{-1}) \geq \text{rmin}\{V_V^+(f(x)), V_V^+(f(y))\}.$$

Similarly,

$$V_V^-(f(x) - f(y)) = V_V^-(f(x - y)) \leq V_A^-(x - y) \leq \text{rmax}\{V_A^-(x), V_A^-(y)\} = \text{rmax}\{V_V^-(f(x)), V_V^-(f(y))\},$$

which implies that

$$V_V^-(f(x) - f(y)) \leq \text{rmax}\{V_V^-(f(x)), V_V^-(f(y))\}.$$

And,

$$V_V^-(f(x)f(y)^{-1}) = V_V^-(f(xy^{-1})) \leq V_A^-(xy^{-1}) \leq \text{rmax}\{V_A^-(x), V_A^-(y)\} = \text{rmax}\{V_V^-(f(x)), V_V^-(f(y))\},$$

which implies that

$$V_V^-(f(x)f(y)^{-1}) \leq \text{rmax}\{V_V^-(f(x)), V_V^-(f(y))\}.$$

Hence V is a $\mathbf{B}_{VV}SF$ of F_2 . □

Theorem 3.2

The homomorphic preimage of a $\mathbf{B}_{VV}SF$ of a field F_2 is a $\mathbf{B}_{VV}SF$ of a field F_1 .

Proof

Let $f : F_1 \rightarrow F_2$ be a homomorphism. Let $V = f(A) = \langle V_V^+, V_V^- \rangle$ be a $\mathbf{B}_{VV}SF$ of F_2 . We have to prove that $A = \langle V_A^+, V_A^- \rangle$ is a $\mathbf{B}_{VV}SF$ of F_1 . Let $x, y \in F_1$. Now

$$V_A^+(x - y) = V_V^+(f(x - y)) = V_V^+(f(x) - f(y)) \geq \text{rmin}\{V_V^+(f(x)), V_V^+(f(y))\} = \text{rmin}\{V_A^+(x), V_A^+(y)\},$$

which implies that

$$V_A^+(x - y) \geq \text{rmin}\{V_A^+(x), V_A^+(y)\}.$$

Also,

$$V_A^+(xy^{-1}) = V_V^+(f(xy^{-1})) = V_V^+(f(x)f(y)^{-1}) \geq \text{rmin}\{V_V^+(f(x)), V_V^+(f(y))\} = \text{rmin}\{V_A^+(x), V_A^+(y)\},$$

which implies that

$$V_A^+(xy^{-1}) \geq \text{rmin}\{V_A^+(x), V_A^+(y)\}.$$

Similarly,

$$V_A^-(x - y) = V_V^-(f(x - y)) = V_V^-(f(x) - f(y)) \leq \text{rmax}\{V_V^-(f(x)), V_V^-(f(y))\} = \text{rmax}\{V_A^-(x), V_A^-(y)\},$$

which implies that

$$V_A^-(x - y) \leq \text{rmax}\{V_A^-(x), V_A^-(y)\}.$$

And,

$$V_A^-(xy^{-1}) = V_V^-(f(xy^{-1})) = V_V^-(f(x)f(y)^{-1}) \leq \text{rmax}\{V_V^-(f(x)), V_V^-(f(y))\} = \text{rmax}\{V_A^-(x), V_A^-(y)\},$$

which implies that

$$V_A^-(xy^{-1}) \leq \text{rmax}\{V_A^-(x), V_A^-(y)\}.$$

Hence A is a $\mathbf{B}_{VV}SF$ of F_1 . □

Theorem 3.3

The homomorphic image of a product of two $\mathbf{B}_{VV}SF$ s of the fields F_1 and F_2 is a $\mathbf{B}_{VV}SF$ of a field.

Proof

From Theorems 2.1 and 3.1, the proof follows directly. □

Theorem 3.4

The homomorphic preimage of a product of two $\mathbf{B}_{VV}SF$ s of the fields F_1 and F_2 is a $\mathbf{B}_{VV}SF$ of a field.

Proof

From Theorems 2.1 and 3.2, the proof follows directly. □

Conclusion

Using the above theorems, we can find more results. This study can be extended to different types of \mathbf{B}_{VV} algebra, particular, \mathbf{B}_{VV} subspaces and \mathbf{B}_{VV} normed spaces are natural extension of the $\mathbf{B}_{VV}SF$ of a field of this work. Field theory has varies types of application in natural life that the application depends the crisp field but after the introduction of fuzzy set and their extension the crisp concept is extended to uncertainty concepts and their extension, the one of the extension is bipolar valued vague subfield of the field. Using these we can develop the application in real field.

REFERENCES

1. B. Anandh and R. Giri, "Notes on Intuitionistic (T, S)-Fuzzy Subfields of A Field," *IOSR Journal of Mathematics (IOSR-JM)*, vol. 12, no. 5, ver. III, pp. 30–34, 2016.
2. M. S. Anitha, M. Muruganatha Prasad, and K. Arjunan, "Notes on bipolar valued fuzzy subgroups of a group," *Bulletin of Society for Mathematical Services and Standards*, vol. 2, no. 3, pp. 52–59, 2013.
3. K. Bala Bavithra, M. Muthusamy, and K. Arjunan, "A Research in Bipolar valued vague subfields of a field," *Communications on Applied Nonlinear Analysis*, vol. 31, no. 7s, pp. 414–420, 2024.
4. A. Balasubramanian, K. L. Muruganatha Prasad, and K. Arjunan, "Properties of Bipolar interval valued fuzzy subgroups of a group," *International Journal of Scientific Research*, vol. 4, no. 4, pp. 262–268, 2015.
5. R. Biswas, "Vague groups," *International Journal of Computational Cognition*, vol. 4, no. 2, pp. 20–23, 2006.

6. S. Cicily Flora and I. Arockiarani, "A new class of generalized bipolar vague sets," *International Journal of Information Research and Review*, vol. 3, no. 11, pp. 3058–3065, 2016.
7. B. Deebea, S. Naganathan, and K. Arjunan, "A study on bipolar valued vague subrings of a ring," *Journal of Shanghai Jiaotong University*, vol. 16, no. 10, pp. 512–518, 2020.
8. B. Deepa, S. Naganathan, and K. Arjunan, "Homomorphism and anti homomorphism functions in Bipolar valued vague subrings of a ring," *International Journal of Mathematical Archive*, vol. 12, no. 7, pp. 1–5, 2021.
9. B. Deebea, S. Naganathan, and K. Arjunan, "A Research on Bipolar valued vague normal subrings of a ring," *Mathematical Statistician and Engineering Applications*, vol. 72, no. 1, pp. 1926–1933, 2023.
10. W. L. Gau and D. J. Buehrer, "Vague sets," *IEEE Transactions on Systems, Man and Cybernetics*, vol. 23, pp. 610–614, 1993.
11. I. Grattan-Guinness, "Fuzzy membership mapped onto interval and many valued quantities," *Z. Math. Logik. Grundlehren Math.*, vol. 22, pp. 149–160, 1975.
12. K. M. Lee, "Bipolar valued fuzzy sets and their operations," in *Proc. Int. Conf. on Intelligent Technologies*, Bangkok, Thailand, 2000, pp. 307–312.
13. K. M. Lee, "Comparison of interval valued fuzzy sets, intuitionistic fuzzy sets and bipolar valued fuzzy sets," *J. Fuzzy Logic Intelligent Systems*, vol. 14, no. 2, pp. 125–129, 2004.
14. M. Muthusamy, N. Palaniappan, and K. Arjunan, "Homomorphism and anti homomorphism of level subfield of intuitionistic fuzzy subfield of a field," *International Journal of Computational and Applied Mathematics*, vol. 4, no. 3, pp. 299–306, 2009.
15. A. Rosenfeld, "Fuzzy groups," *Journal of Mathematical Analysis and Applications*, vol. 35, pp. 512–517, 1971.
16. C. Yamini, K. Arjunan, and B. Ananth, "Bipolar valued multi fuzzy subfield of a field," *International Journal of Management, Technology And Engineering*, vol. 8, no. 11, pp. 1706–1711, 2018.
17. B. Yasodara and K. E. Sathappan, "Bipolar-valued multi fuzzy subsemifields of a semifield," *International Journal of Mathematical Archive*, vol. 6, no. 9, pp. 75–80, 2015.
18. L. A. Zadeh, "Fuzzy sets," *Information and Control*, vol. 8, pp. 338–353, 1965.
19. W. R. Zhang, "Bipolar Fuzzy sets and Relations, a computational framework for cognitive modeling and multiple decision analysis," in *Proc. IEEE International Conference on Fuzzy Systems*, 1994, pp. 305–309.